

National Core Indicators[®] - Intellectual and Developmental Disabilities (NCI[®]-IDD)



National Report 2022-23

Health




Table of Contents: Health

| | |
|--|----------|
| Where do these data come from? | 4 |
| What are “collapsed” data? | 4 |
| What is weighting and why do we do it? | 4 |
| Presentation of state and weighted NCI-IDD averages | 5 |
| Averages..... | 5 |
| Important Note on Changes to the 2022-23 Survey Tool | 5 |
| Charts and Tables | 7 |
| Table 1. Has a primary care doctor or practitioner | 9 |
| Table 1b. Percent who have a primary care doctor or practitioner (includes missing and “don’t know” responses)..... | 10 |
| Table 2. In poor health..... | 12 |
| Table 3. Had a complete physical exam in the past year (routine exam, not a visit for a specific problem or illness)..... | 14 |
| Table 3b. Last physical exam (includes missing and “don’t know” responses)..... | 15 |
| Table 4. Had a routine dental exam in the past year..... | 17 |
| Table 4b. Last dental exam (includes missing and “don’t know” responses) | 18 |
| Table 5. Had an eye exam/vision screening in the past year..... | 20 |
| Table 5b. Last eye exam or vision screening (includes missing and “don’t know” responses) | 21 |
| Table 6. Had a hearing test in the past five years..... | 23 |
| Table 6b. Had a hearing test in the past five years (includes missing and “don’t know” responses)..... | 24 |
| Table 7. Had a Pap test in the past three years (among women 21 and older) | 26 |
| Table 7b. Last Pap test (among women 21 and older, includes “not applicable,” missing, and “don’t know” responses) | 27 |
| Table 8. Had a mammogram test in the past two years (among women age 50 and over) | 29 |
| Table 8b. Last mammogram test (among women age 40 and over, includes “not applicable,” missing, and “don’t know”) | 30 |
| Table 9. Last colorectal cancer screening (among people age 45 to 75)..... | 32 |
| Table 9b. Last colorectal cancer screening (among people age 45 to 75, includes “not applicable,” missing, and “don’t know” responses) | 33 |
| Table 10. Had a flu vaccine in the past 12 months..... | 35 |
| Table 10b. Had a flu vaccine in the past year (includes missing and “don’t know” responses)..... | 36 |
| Table 11. Went to the emergency room for any reason to get care for themselves in the past 12 months | 38 |

| | |
|---|----|
| Table 12. Had an unexpected hospital admission due to dehydration, bowel obstruction, seizure, aspiration, or GERD, in the past year..... | 40 |
| Table 13. Fell and hurt themselves in the past six months | 42 |

Where do these data come from?

Most items in this section are drawn from administrative sources. Select items may have been collected during the face-to-face survey from the person receiving services, or for some select questions, from a “proxy” respondent. A proxy respondent is someone who knows the person well (e.g., a family member), proxy respondents may not be a case manager/service coordinator. Items that may come from administrative records or allow for proxy respondents are noted in the table title. Read more about why we allow proxy respondents for select questions in the IPS in the Overview chapter of this report.

Items in this chapter are a part of the NCI-IDD Individuals Outcomes domain (or topic area) and comprise the “Health” subdomain (a specific area of concern within the domain). The Health subdomain includes items that help determine whether “People have access to and get recommended health services at the recommended frequencies”.

Read more about NCI-IDD domains and sub-domains in the Overview chapter of this report.

What are “collapsed” data?

Many questions in the IPS allow respondents to answer a question with “middle” response like “maybe”, “sometimes” or “in-between”. The tables in this section present collapsed data. That means that two or more response options are grouped together if they are considered to reflect a positive outcome.

To see the breakout of responses for each option, please see Appendix C.

What is weighting and why do we do it?

The NCI-IDD averages are “weighted” to reflect the states’ relative population and sample sizes. We created the weights using the state’s number of valid surveys and its total survey-eligible population. This way, a state that provides services to a larger number of people but uses a sample similar in size to other states has a greater influence on the overall NCI-IDD average (that is, its contribution is *proportional to its service population*). You can read more about this in the Methodology section of the Overview chapter.

The number of respondents (N) for each question *is not* weighted.

Presentation of state and weighted NCI-IDD averages

Averages

For the most part, the state outcomes are shown in descending order of percentage and are categorized as statistically significantly above the NCI-IDD Average, within the range of the NCI-IDD Average, or significantly below the NCI-IDD Average. Statistical significance is taken to be at or below the .01 level. For those states that fall within the NCI-IDD Average range, their 'always' or 'yes' response was not statistically different from the NCI-IDD Average. Other tables show the states listed in alphabetical order and are not tested for significance.

Tables that show significance also break out average results for each state by the type of residence: ICF/IID or other institutional settings, group residence, own home, parent/relative's home, foster or host home. Figure A on the following page shows, by state, the percentage of people surveyed in each residence type. Please note though that:

- Significance testing was not performed on the breakouts by residence
- The number of people in each residential setting is often too small to allow for valid state-to-state comparisons
- And in many cases, statistically valid conclusions cannot be drawn about differences in results between residence types

Still, we provide this information for states' internal purposes. It should not be used to compare one state with another.

To protect people's privacy, we do not show the data when 20 or fewer people from a state or residence type responded to a question (though their responses are included in the NCI-IDD average).

Important Note on Changes to the 2022-23 Survey Tool

NCI-IDD operates on a six-year revisions cycle to regularly ensure the current set of indicators meet the needs of state systems and reflect current needs in the field. Survey changes are guided by state managers, people with disabilities, and other experts in the field. The 2021-22 survey tool reflected changes based on this guidance and additional survey testing. Please note, any changes to the survey tool may affect comparison to previous years' data, therefore any cross-year comparison should be made with caution.

Figure A. Percentage of people surveyed in each residence type by state

| State (names abbreviated) | Institutional settings | Community-based Group Residential settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care/Host Home | N |
|---------------------------------|------------------------|--|-----------------------|-----------------------------|-----------------------|-------|
| AL | 0% | 69% | 10% | 21% | 0% | 477 |
| AR | 0% | 13% | 43% | 41% | 4% | 417 |
| CA | 7% | 23% | 13% | 56% | 1% | 8808 |
| CT | 0% | 48% | 14% | 35% | 3% | 600 |
| DE | 0% | 54% | 3% | 40% | 3% | 414 |
| DC | 18% | 38% | 34% | 6% | 5% | 133 |
| FL | 0% | 36% | 14% | 50% | 0% | 524 |
| GA | 0% | 37% | 12% | 39% | 12% | 406 |
| HI | 0% | 4% | 3% | 52% | 41% | 344 |
| IL | 2% | 47% | 7% | 43% | 1% | 398 |
| IN | 31% | 19% | 9% | 41% | 0% | 1074 |
| KS | 0% | 48% | 20% | 30% | 3% | 312 |
| KY | 0% | 28% | 9% | 40% | 23% | 437 |
| MD | 1% | 32% | 10% | 57% | 0% | 522 |
| MI | 0% | 37% | 18% | 41% | 3% | 652 |
| MN | 0% | 47% | 9% | 39% | 4% | 401 |
| MO | 2% | 24% | 33% | 39% | 3% | 400 |
| MT | 0% | 48% | 31% | 20% | 0% | 365 |
| NE | 5% | 25% | 15% | 25% | 30% | 420 |
| NV | 0% | 48% | 14% | 32% | 6% | 416 |
| NH | 1% | 15% | 9% | 41% | 33% | 392 |
| NJ | 0% | 38% | 13% | 47% | 2% | 459 |
| NY | 0% | 41% | 7% | 51% | 1% | 1662 |
| NC | 6% | 21% | 6% | 51% | 15% | 590 |
| OK | 0% | 45% | 19% | 35% | 2% | 400 |
| OR | 0% | 15% | 10% | 61% | 14% | 418 |
| PA | 8% | 38% | 11% | 39% | 4% | 663 |
| SC | 1% | 51% | 14% | 29% | 5% | 509 |
| TX | 5% | 7% | 8% | 39% | 40% | 416 |
| UT | 7% | 40% | 10% | 28% | 16% | 352 |
| VA | 0% | 36% | 9% | 39% | 17% | 757 |
| WI | 4% | 26% | 31% | 39% | 0% | 321 |
| WY | 0% | 0% | 45% | 55% | 0% | 269 |
| Weighted NCI-IDD Average | 4% | 31% | 13% | 46% | 6% | 24728 |

Table note: Institutional settings includes ICF/IID, Nursing Facilities, and other institutional settings

Charts and Tables

Person has primary care doctor/practitioner




Table 1. Has a primary care doctor or practitioner*Information may have been obtained through state records*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| DC * ↑ | 100% | 97 | n/a | 100% | 100% | n/a | n/a |
| DE ↑ | 100% | 391 | n/a | 100% | n/a | 100% | n/a |
| IN ↑ | 100% | 1,080 | 100% | 100% | 100% | 100% | n/a |
| HI ↑ | 99% | 350 | n/a | n/a | n/a | 100% | 99% |
| KS ↑ | 99% | 314 | n/a | 99% | 100% | 100% | n/a |
| GA ↑ | 99% | 406 | n/a | 100% | 98% | 99% | 100% |
| NY ↑ | 99% | 1,674 | n/a | 99% | 99% | 99% | n/a |
| AL ↑ | 99% | 469 | n/a | 99% | 100% | 98% | n/a |
| AR ↑ | 99% | 414 | n/a | 98% | 98% | 99% | n/a |
| VA ↑ | 98% | 726 | n/a | 100% | 98% | 97% | 98% |
| OK | 99% | 400 | n/a | 99% | 100% | 97% | n/a |
| MI | 98% | 660 | n/a | 99% | 96% | 98% | 100% |
| NH | 98% | 395 | n/a | 100% | 92% | 98% | 98% |
| CT | 98% | 600 | n/a | 99% | 98% | 97% | n/a |
| MO | 98% | 403 | n/a | 99% | 98% | 97% | n/a |
| NE | 98% | 416 | 100% | 99% | 93% | 98% | 98% |
| MD | 98% | 522 | n/a | 99% | 96% | 97% | n/a |
| PA | 97% | 664 | 100% | 95% | 99% | 98% | 100% |
| IL | 97% | 387 | n/a | 98% | 96% | 96% | n/a |
| UT | 97% | 351 | 96% | 98% | 94% | 97% | 98% |
| NCI-IDD Average | 97% | 24,572 | 99% | 98% | 96% | 96% | 97% |
| NV | 97% | 393 | n/a | 98% | 98% | 93% | 100% |
| NC | 97% | 589 | 100% | 98% | 97% | 95% | 100% |
| NJ | 97% | 454 | n/a | 97% | 97% | 97% | n/a |
| KY | 97% | 431 | n/a | 97% | 100% | 96% | 96% |
| OR | 97% | 401 | n/a | 100% | 98% | 95% | 100% |
| SC | 96% | 510 | n/a | 99% | 93% | 95% | 100% |
| MN | 96% | 386 | n/a | 98% | 97% | 93% | n/a |
| WI * ↓ | 96% | 254 | n/a | 96% | 95% | 96% | n/a |
| FL | 96% | 528 | n/a | 97% | 92% | 96% | n/a |
| MT | 95% | 361 | n/a | 95% | 95% | 94% | n/a |
| TX | 94% | 390 | 100% | 93% | 97% | 93% | 94% |
| WY ↓ | 67% | 424 | n/a | n/a | 67% | 66% | n/a |

Table note: States with more than 25% missing or don't know are marked with *

Table 1b. Percent who have a primary care doctor or practitioner (includes missing and “don’t know” responses)

| State (names abbreviated) | Has a primary care doctor or practitioner | Does not have primary care doctor or practitioner | Missing | Don't Know | N |
|---------------------------|---|---|-----------|------------|---------------|
| AL | 97% | 1% | 2% | 0% | 479 |
| AR | 98% | 1% | 1% | 0% | 417 |
| CT | 98% | 2% | 0% | 0% | 602 |
| DE | 93% | 0% | 0% | 7% | 419 |
| DC * | 27% | 0% | 63% | 10% | 354 |
| FL | 95% | 4% | 1% | 0% | 532 |
| GA | 99% | 1% | 0% | 0% | 406 |
| HI | 99% | 1% | 0% | 0% | 351 |
| IL | 94% | 3% | 2% | 2% | 400 |
| IN | 100% | 0% | 0% | 0% | 1,080 |
| KS | 79% | 1% | 18% | 2% | 393 |
| KY | 95% | 3% | 1% | 0% | 439 |
| MD | 95% | 2% | 0% | 2% | 533 |
| MI | 97% | 2% | 0% | 1% | 670 |
| MN | 92% | 3% | 4% | 0% | 403 |
| MO | 98% | 2% | 0% | 0% | 403 |
| MT | 94% | 5% | 1% | 0% | 365 |
| NE | 97% | 2% | 0% | 0% | 420 |
| NV | 91% | 3% | 1% | 5% | 419 |
| NH | 97% | 2% | 1% | 0% | 399 |
| NJ | 96% | 3% | 1% | 0% | 460 |
| NY | 98% | 1% | 0% | 0% | 1,682 |
| NC | 96% | 3% | 1% | 0% | 594 |
| OK | 99% | 2% | 0% | 0% | 400 |
| OR | 93% | 3% | 1% | 3% | 418 |
| PA | 90% | 3% | 6% | 2% | 718 |
| SC | 96% | 4% | 0% | 0% | 511 |
| TX | 87% | 6% | 7% | 0% | 420 |
| UT | 95% | 3% | 1% | 1% | 359 |
| VA | 94% | 1% | 1% | 4% | 762 |
| WI * | 67% | 3% | 29% | 1% | 362 |
| WY | 67% | 33% | 0% | 0% | 424 |
| NCI-IDD Average | 93% | 3% | 3% | 1% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Overall, how would you describe your health?




Table 2. In poor health*Proxy respondents were allowed for this question*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| NE ↑ | 6% | 399 | n/a | 8% | 3% | 2% | 10% |
| WI | 5% | 351 | n/a | 1% | 10% | 1% | n/a |
| OR | 4% | 404 | n/a | 5% | 10% | 2% | 9% |
| KY | 4% | 433 | n/a | 4% | 0% | 5% | 5% |
| KS | 3% | 377 | n/a | 2% | 5% | 1% | n/a |
| AR | 3% | 412 | n/a | 6% | 1% | 4% | n/a |
| MT | 3% | 348 | n/a | 3% | 3% | 1% | n/a |
| WY | 3% | 405 | n/a | n/a | 4% | 1% | n/a |
| DE | 3% | 410 | n/a | 4% | n/a | 1% | n/a |
| FL | 3% | 525 | n/a | 3% | 3% | 2% | n/a |
| TX | 3% | 397 | 0% | 0% | 3% | 2% | 4% |
| VA | 2% | 760 | n/a | 4% | 2% | 2% | 1% |
| OK | 2% | 397 | n/a | 3% | 3% | 1% | n/a |
| SC | 2% | 504 | n/a | 3% | 1% | 1% | 0% |
| NCI-IDD Average | 2% | 24,920 | 2% | 2% | 3% | 2% | 3% |
| MN | 2% | 390 | n/a | 1% | 3% | 3% | n/a |
| NV | 2% | 403 | n/a | 2% | 4% | 1% | 4% |
| NY | 2% | 1,618 | n/a | 3% | 1% | 1% | n/a |
| NJ | 2% | 450 | n/a | 2% | 2% | 1% | n/a |
| MD | 2% | 513 | n/a | 2% | 2% | 2% | n/a |
| CT | 2% | 584 | n/a | 1% | 2% | 1% | n/a |
| NC | 2% | 585 | 0% | 0% | 0% | 2% | 2% |
| UT | 2% | 326 | 0% | 3% | 3% | 0% | 0% |
| MI | 1% | 663 | n/a | 2% | 2% | 1% | 0% |
| PA | 1% | 692 | 0% | 2% | 0% | 1% | 4% |
| NH | 1% | 398 | n/a | 0% | 3% | 2% | 1% |
| HI | 1% | 348 | n/a | n/a | n/a | 1% | 0% |
| GA | 1% | 405 | n/a | 2% | 0% | 0% | 2% |
| DC | 1% | 346 | 0% | 0% | 2% | n/a | n/a |
| AL ↓ | 1% | 470 | n/a | 1% | 0% | 1% | n/a |
| IN ↓ | 1% | 1,070 | 1% | 1% | 0% | 1% | n/a |
| IL ↓ | 1% | 388 | n/a | 0% | 4% | 0% | n/a |
| MO ↓ | 1% | 399 | n/a | 0% | 2% | 0% | n/a |

Last complete annual physical exam




Table 3. Had a complete physical exam in the past year (routine exam, not a visit for a specific problem or illness)

Information may have been obtained through state records

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------|----------------------|---------------|------------------------|----------------|-----------------------|-----------------------------|--------------------------|
| NJ ↑ | 98% | 444 | n/a | 99% | 100% | 96% | n/a |
| KY ↑ | 97% | 414 | n/a | 98% | 100% | 95% | 96% |
| GA ↑ | 96% | 398 | n/a | 98% | 93% | 96% | 94% |
| OK ↑ | 96% | 400 | n/a | 96% | 95% | 96% | n/a |
| AR ↑ | 95% | 398 | n/a | 92% | 95% | 96% | n/a |
| AL ↑ | 95% | 456 | n/a | 97% | 94% | 90% | n/a |
| CT ↑ | 95% | 596 | n/a | 97% | 98% | 92% | n/a |
| FL ↑ | 95% | 525 | n/a | 98% | 93% | 93% | n/a |
| NH ↑ | 94% | 383 | n/a | 98% | 94% | 91% | 97% |
| MI ↑ | 94% | 507 | n/a | 97% | 89% | 93% | n/a |
| SC ↑ | 93% | 481 | n/a | 96% | 92% | 87% | 100% |
| PA ↑ | 93% | 627 | 96% | 96% | 94% | 90% | 91% |
| NE ↑ | 93% | 416 | n/a | 95% | 97% | 86% | 95% |
| IL ↑ | 92% | 330 | n/a | 97% | 91% | 85% | n/a |
| MO ↑ | 92% | 395 | n/a | 96% | 94% | 90% | n/a |
| UT ↑ | 92% | 311 | 95% | 95% | 93% | 88% | 96% |
| MT ↑ | 92% | 353 | n/a | 93% | 94% | 87% | n/a |
| IN ↑ | 92% | 1,034 | 95% | 92% | 95% | 88% | n/a |
| MD ↑ | 90% | 495 | n/a | 92% | 86% | 89% | n/a |
| TX ↑ | 88% | 382 | n/a | 93% | 91% | 84% | 92% |
| NC ↑ | 88% | 544 | 100% | 92% | 85% | 84% | 91% |
| MN ↑ | 88% | 382 | n/a | 92% | 89% | 81% | n/a |
| NCI-IDD Average | 83% | 23,463 | 94% | 86% | 84% | 79% | 91% |
| WY * | 82% | 284 | n/a | n/a | 86% | 81% | n/a |
| WI * | 81% | 241 | n/a | 82% | 81% | 83% | n/a |
| HI | 81% | 342 | n/a | n/a | n/a | 80% | 82% |
| KS | 80% | 305 | n/a | 84% | 78% | 76% | n/a |
| OR | 80% | 379 | n/a | 93% | 68% | 75% | 95% |
| DC * | 77% | 133 | 74% | 90% | 77% | n/a | n/a |
| VA ↓ | 78% | 743 | n/a | 82% | 79% | 75% | 78% |
| NV ↓ | 73% | 373 | n/a | 74% | 67% | 74% | 75% |
| DE * ↓ | 60% | 238 | n/a | 66% | n/a | 51% | n/a |
| NY ↓ | 53% | 1,609 | n/a | 55% | 46% | 52% | n/a |

Table note: States with more than 25% missing or don't know are marked with *

Table 3b. Last physical exam (includes missing and “don’t know” responses)

| State (names abbreviated) | In the past year | More than 1 year | Missing | Don't Know | N |
|---------------------------|------------------|------------------|-----------|------------|---------------|
| AL | 91% | 4% | 2% | 3% | 479 |
| AR | 91% | 4% | 1% | 4% | 417 |
| CT | 94% | 5% | 0% | 1% | 602 |
| DE * | 34% | 23% | 0% | 43% | 419 |
| DC * | 29% | 9% | 62% | 0% | 354 |
| FL | 94% | 5% | 1% | 1% | 532 |
| GA | 94% | 4% | 0% | 2% | 406 |
| HI | 79% | 19% | 2% | 1% | 351 |
| IL | 76% | 6% | 3% | 14% | 400 |
| IN | 88% | 8% | 0% | 4% | 1,080 |
| KS | 62% | 15% | 19% | 4% | 393 |
| KY | 91% | 3% | 1% | 5% | 439 |
| MD | 84% | 9% | 1% | 6% | 533 |
| MI | 71% | 5% | 0% | 24% | 670 |
| MN | 83% | 12% | 4% | 1% | 403 |
| MO | 91% | 7% | 0% | 2% | 403 |
| MT | 89% | 8% | 1% | 3% | 365 |
| NE | 92% | 7% | 0% | 0% | 420 |
| NV | 65% | 24% | 2% | 9% | 419 |
| NH | 90% | 6% | 2% | 2% | 399 |
| NJ | 94% | 2% | 1% | 2% | 460 |
| NY | 51% | 45% | 0% | 4% | 1,682 |
| NC | 81% | 11% | 1% | 8% | 594 |
| OK | 96% | 4% | 0% | 0% | 400 |
| OR | 72% | 18% | 1% | 8% | 418 |
| PA | 81% | 6% | 7% | 6% | 718 |
| SC | 88% | 6% | 0% | 6% | 511 |
| TX | 80% | 10% | 7% | 2% | 420 |
| UT | 80% | 7% | 3% | 10% | 359 |
| VA | 76% | 21% | 1% | 1% | 762 |
| WI * | 54% | 12% | 29% | 4% | 362 |
| WY * | 55% | 12% | 0% | 33% | 424 |
| NCI-IDD Average | 76% | 15% | 3% | 5% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last dentist exam




Table 4. Had a routine dental exam in the past year*Information may have been obtained through state records*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| NE ↑ | 91% | 418 | 100% | 90% | 95% | 87% | 94% |
| NJ ↑ | 89% | 422 | n/a | 98% | 87% | 82% | n/a |
| CT ↑ | 88% | 586 | n/a | 93% | 85% | 82% | n/a |
| AL ↑ | 87% | 432 | n/a | 92% | 82% | 68% | n/a |
| IN ↑ | 86% | 1,028 | 92% | 88% | 84% | 81% | n/a |
| KY ↑ | 86% | 381 | n/a | 92% | 87% | 77% | 89% |
| SC ↑ | 85% | 453 | n/a | 94% | 79% | 67% | 87% |
| UT ↑ | 85% | 273 | 95% | 90% | 79% | 73% | 93% |
| PA ↑ | 84% | 583 | 88% | 95% | 70% | 76% | 81% |
| MN ↑ | 84% | 377 | n/a | 86% | 75% | 81% | n/a |
| MT ↑ | 83% | 346 | n/a | 82% | 88% | 79% | n/a |
| IL | 81% | 313 | n/a | 84% | 76% | 74% | n/a |
| GA | 79% | 387 | n/a | 86% | 71% | 73% | 83% |
| DE * | 79% | 207 | n/a | 80% | n/a | 76% | n/a |
| TX | 79% | 383 | n/a | 89% | 74% | 72% | 83% |
| MO | 79% | 382 | n/a | 89% | 83% | 68% | n/a |
| NH | 78% | 346 | n/a | 91% | 68% | 79% | 73% |
| DC * | 78% | 127 | 77% | 85% | 83% | n/a | n/a |
| MD | 78% | 489 | n/a | 86% | 75% | 74% | n/a |
| OK | 77% | 396 | n/a | 82% | 72% | 72% | n/a |
| FL | 77% | 509 | n/a | 87% | 89% | 66% | n/a |
| NCI-IDD Average | 76% | 22,553 | 90% | 83% | 73% | 70% | 81% |
| KS * | 75% | 259 | n/a | 78% | 79% | 69% | n/a |
| NC | 75% | 493 | 100% | 76% | 70% | 69% | 79% |
| MI * | 72% | 434 | n/a | 72% | 69% | 74% | n/a |
| AR | 70% | 318 | n/a | 77% | 73% | 67% | n/a |
| OR ↓ | 70% | 386 | n/a | 68% | 62% | 71% | 75% |
| WI * ↓ | 67% | 223 | n/a | 67% | 60% | 72% | n/a |
| WY ↓ | 66% | 320 | n/a | n/a | 62% | 68% | n/a |
| NY ↓ | 62% | 1,480 | n/a | 71% | 60% | 55% | n/a |
| VA ↓ | 59% | 724 | n/a | 59% | 63% | 55% | 68% |
| NV * ↓ | 56% | 277 | n/a | 56% | 51% | 58% | n/a |
| HI ↓ | 52% | 269 | n/a | n/a | n/a | 50% | 60% |

*Table note: States with more than 25% missing or don't know are marked with **

Table 4b. Last dental exam (includes missing and “don’t know” responses)

| State (names abbreviated) | More than 1 year | In the past year | Missing | Don't Know | N |
|---------------------------|------------------|------------------|-----------|------------|---------------|
| AL | 12% | 78% | 2% | 8% | 479 |
| AR | 23% | 53% | 1% | 23% | 417 |
| CT | 12% | 85% | 0% | 2% | 602 |
| DE * | 11% | 39% | 0% | 51% | 419 |
| DC * | 8% | 28% | 63% | 1% | 354 |
| FL | 22% | 74% | 1% | 3% | 532 |
| GA | 20% | 75% | 0% | 5% | 406 |
| HI | 36% | 40% | 13% | 11% | 351 |
| IL | 15% | 63% | 2% | 20% | 400 |
| IN | 13% | 82% | 0% | 5% | 1,080 |
| KS * | 17% | 49% | 19% | 15% | 393 |
| KY | 13% | 74% | 1% | 13% | 439 |
| MD | 20% | 71% | 1% | 8% | 533 |
| MI * | 18% | 47% | 0% | 35% | 670 |
| MN | 15% | 78% | 4% | 3% | 403 |
| MO | 20% | 74% | 0% | 5% | 403 |
| MT | 16% | 79% | 1% | 5% | 365 |
| NE | 9% | 91% | 0% | 0% | 420 |
| NV * | 29% | 37% | 2% | 32% | 419 |
| NH | 19% | 68% | 2% | 12% | 399 |
| NJ | 10% | 82% | 2% | 6% | 460 |
| NY | 33% | 55% | 0% | 12% | 1,682 |
| NC | 21% | 62% | 1% | 16% | 594 |
| OK | 23% | 76% | 0% | 1% | 400 |
| OR | 28% | 64% | 1% | 6% | 418 |
| PA | 13% | 68% | 6% | 13% | 718 |
| SC | 14% | 75% | 0% | 11% | 511 |
| TX | 20% | 72% | 7% | 2% | 420 |
| UT | 12% | 64% | 3% | 21% | 359 |
| VA | 39% | 56% | 1% | 4% | 762 |
| WI * | 20% | 41% | 29% | 10% | 362 |
| WY | 25% | 50% | 0% | 25% | 424 |
| NCI-IDD Average | 21% | 67% | 3% | 9% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last eye exam/vision screening




Table 5. Had an eye exam/vision screening in the past year*Information may have been obtained through state records*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| GA ↑ | 79% | 363 | n/a | 86% | 77% | 71% | 80% |
| DC * ↑ | 78% | 96 | n/a | 81% | 75% | n/a | n/a |
| IL * ↑ | 77% | 285 | n/a | 89% | 67% | 51% | n/a |
| SC ↑ | 73% | 439 | n/a | 90% | 61% | 38% | n/a |
| NJ ↑ | 71% | 384 | n/a | 86% | 75% | 55% | n/a |
| KS * ↑ | 68% | 238 | n/a | 76% | 67% | 51% | n/a |
| HI ↑ | 68% | 299 | n/a | n/a | n/a | 70% | 67% |
| KY * ↑ | 67% | 309 | n/a | 80% | 65% | 51% | 74% |
| PA * ↑ | 66% | 535 | 71% | 83% | 55% | 47% | 82% |
| OK ↑ | 65% | 393 | n/a | 78% | 75% | 44% | n/a |
| MN ↑ | 64% | 365 | n/a | 77% | 51% | 52% | n/a |
| MD ↑ | 62% | 448 | n/a | 73% | 67% | 56% | n/a |
| AL ↑ | 61% | 401 | n/a | 64% | 59% | 51% | n/a |
| AR | 59% | 330 | n/a | 63% | 71% | 49% | n/a |
| DE * | 58% | 93 | n/a | 72% | n/a | 14% | n/a |
| CT | 58% | 559 | n/a | 68% | 67% | 36% | n/a |
| IN | 55% | 1,005 | 58% | 61% | 54% | 51% | n/a |
| NE | 55% | 417 | 65% | 63% | 59% | 46% | 52% |
| UT * | 55% | 164 | 82% | 51% | n/a | 30% | 77% |
| MO | 55% | 373 | n/a | 61% | 63% | 43% | n/a |
| NCI-IDD Average | 53% | 20,921 | 64% | 62% | 54% | 45% | 59% |
| NH | 53% | 325 | n/a | 67% | 60% | 40% | 57% |
| VA * | 51% | 325 | n/a | 52% | 48% | 44% | 68% |
| NC * | 49% | 390 | 61% | 59% | 32% | 40% | 58% |
| WI * | 47% | 215 | n/a | 42% | 59% | 40% | n/a |
| OR | 47% | 365 | n/a | 51% | 49% | 42% | 67% |
| WY ↓ | 45% | 319 | n/a | n/a | 42% | 47% | n/a |
| FL ↓ | 45% | 482 | n/a | 60% | 51% | 33% | n/a |
| MI * ↓ | 43% | 343 | n/a | 36% | 51% | 43% | n/a |
| MT ↓ | 42% | 327 | n/a | 41% | 43% | 44% | n/a |
| TX ↓ | 42% | 371 | n/a | 36% | 40% | 35% | 51% |
| NV * ↓ | 41% | 196 | n/a | 42% | 38% | 49% | n/a |
| NY ↓ | 41% | 1,490 | n/a | 43% | 22% | 41% | n/a |

*Table note: States with more than 25% missing or don't know are marked with **

Table 5b. Last eye exam or vision screening (includes missing and “don’t know” responses)

| State (names abbreviated) | More than 1 year | In the past year | Missing | Don't know | N |
|---------------------------|------------------|------------------|-----------|------------|---------------|
| AL | 33% | 51% | 2% | 14% | 479 |
| AR | 32% | 47% | 0% | 21% | 417 |
| CT | 39% | 54% | 0% | 7% | 602 |
| DE * | 9% | 13% | 0% | 78% | 419 |
| DC * | 6% | 21% | 62% | 11% | 354 |
| FL | 49% | 41% | 2% | 8% | 532 |
| GA | 19% | 70% | 0% | 11% | 406 |
| HI | 27% | 58% | 5% | 9% | 351 |
| IL * | 17% | 55% | 3% | 26% | 400 |
| IN | 42% | 51% | 0% | 7% | 1,080 |
| KS * | 19% | 41% | 19% | 21% | 393 |
| KY * | 23% | 47% | 0% | 29% | 439 |
| MD | 32% | 52% | 1% | 15% | 533 |
| MI * | 29% | 22% | 0% | 49% | 670 |
| MN | 33% | 58% | 4% | 6% | 403 |
| MO | 42% | 51% | 0% | 7% | 403 |
| MT | 52% | 38% | 1% | 10% | 365 |
| NE | 45% | 55% | 0% | 0% | 420 |
| NV * | 27% | 19% | 1% | 53% | 419 |
| NH | 39% | 43% | 1% | 18% | 399 |
| NJ | 24% | 59% | 2% | 14% | 460 |
| NY | 53% | 36% | 0% | 11% | 1,682 |
| NC * | 34% | 32% | 0% | 34% | 594 |
| OK | 34% | 64% | 0% | 2% | 400 |
| OR | 46% | 41% | 1% | 11% | 418 |
| PA * | 25% | 49% | 6% | 20% | 718 |
| SC | 23% | 63% | 0% | 14% | 511 |
| TX | 52% | 37% | 7% | 5% | 420 |
| UT * | 21% | 25% | 3% | 51% | 359 |
| VA * | 21% | 22% | 1% | 57% | 762 |
| WI * | 31% | 28% | 29% | 12% | 362 |
| WY | 41% | 34% | 0% | 25% | 424 |
| NCI-IDD Average | 39% | 44% | 3% | 14% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last hearing test




Table 6. Had a hearing test in the past five years*Information may have been obtained through state records*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| DC * ↑ | 100% | 66 | n/a | n/a | 100% | n/a | n/a |
| PA * ↑ | 85% | 338 | 88% | 94% | 56% | 76% | n/a |
| HI ↑ | 85% | 283 | n/a | n/a | n/a | 86% | 84% |
| UT * ↑ | 63% | 92 | 100% | 53% | n/a | 40% | n/a |
| IL * ↑ | 62% | 198 | n/a | 70% | n/a | 44% | n/a |
| CT ↑ | 59% | 478 | n/a | 74% | 59% | 35% | n/a |
| NE ↑ | 56% | 393 | 80% | 73% | 35% | 50% | 53% |
| IN ↑ | 53% | 938 | 86% | 39% | 40% | 36% | n/a |
| NJ * | 51% | 315 | n/a | 60% | 59% | 42% | n/a |
| AL * | 51% | 316 | n/a | 56% | 53% | 36% | n/a |
| KY * | 50% | 159 | n/a | 63% | n/a | 34% | 67% |
| NY * | 50% | 995 | n/a | 64% | 23% | 44% | n/a |
| KS * | 48% | 147 | n/a | 54% | 30% | 43% | n/a |
| AR * | 47% | 237 | n/a | 57% | 54% | 39% | n/a |
| NCI-IDD Average | 47% | 17,385 | 81% | 58% | 39% | 41% | 38% |
| SC * | 47% | 282 | n/a | 68% | 35% | 27% | n/a |
| NC * | 46% | 272 | 57% | 46% | n/a | 41% | 59% |
| WY * | 45% | 317 | n/a | n/a | 41% | 45% | n/a |
| MI * | 44% | 217 | n/a | 45% | 43% | 43% | n/a |
| MT * | 44% | 219 | n/a | 52% | 41% | 31% | n/a |
| NV * | 44% | 73 | n/a | 42% | n/a | n/a | n/a |
| MN | 43% | 308 | n/a | 55% | 45% | 33% | n/a |
| MD * | 42% | 354 | n/a | 63% | 37% | 35% | n/a |
| NH * | 41% | 213 | n/a | 51% | 24% | 41% | 40% |
| VA * | 40% | 230 | n/a | 43% | 54% | 32% | 50% |
| MO * | 40% | 270 | n/a | 37% | 52% | 27% | n/a |
| FL * ↓ | 36% | 395 | n/a | 42% | 39% | 32% | n/a |
| WI * ↓ | 33% | 105 | n/a | 44% | 21% | 38% | n/a |
| OK ↓ | 30% | 366 | n/a | 36% | 24% | 27% | n/a |
| GA ↓ | 28% | 344 | n/a | 31% | 30% | 22% | 32% |
| TX ↓ | 28% | 337 | n/a | 45% | 19% | 24% | 27% |
| OR ↓ | 21% | 322 | n/a | 22% | 16% | 23% | 15% |

Table note: States with more than 25% missing or don't know are marked with *; The following states had a low count of responses (N<20) and were not shown: DE

Table 6b. Had a hearing test in the past five years (includes missing and “don’t know” responses)

| State (names abbreviated) | Not in past 5 years | Within past 5 years | Missing | Don't know | N |
|---------------------------|---------------------|---------------------|-----------|------------|---------------|
| AL * | 32% | 34% | 3% | 32% | 479 |
| AR * | 30% | 27% | 1% | 42% | 417 |
| CT | 33% | 47% | 0% | 20% | 602 |
| DE * | 2% | 2% | 0% | 96% | 419 |
| DC * | 0% | 19% | 63% | 18% | 354 |
| FL * | 47% | 27% | 2% | 24% | 532 |
| GA | 61% | 23% | 0% | 15% | 406 |
| HI | 12% | 68% | 7% | 13% | 351 |
| IL * | 19% | 31% | 3% | 47% | 400 |
| IN | 41% | 46% | 0% | 13% | 1,080 |
| KS * | 20% | 18% | 19% | 44% | 393 |
| KY * | 18% | 18% | 0% | 64% | 439 |
| MD * | 38% | 28% | 0% | 34% | 533 |
| MI * | 18% | 14% | 1% | 67% | 670 |
| MN | 43% | 33% | 4% | 20% | 403 |
| MO * | 40% | 27% | 0% | 33% | 403 |
| MT * | 34% | 26% | 2% | 38% | 365 |
| NE | 41% | 53% | 1% | 5% | 420 |
| NV * | 10% | 8% | 2% | 81% | 419 |
| NH * | 32% | 22% | 2% | 45% | 399 |
| NJ * | 33% | 35% | 3% | 29% | 460 |
| NY * | 29% | 30% | 0% | 41% | 1,682 |
| NC * | 25% | 21% | 1% | 53% | 594 |
| OK | 64% | 28% | 0% | 9% | 400 |
| OR | 61% | 16% | 1% | 22% | 418 |
| PA * | 7% | 40% | 6% | 47% | 718 |
| SC * | 29% | 26% | 1% | 44% | 511 |
| TX | 58% | 22% | 7% | 13% | 420 |
| UT * | 9% | 16% | 3% | 71% | 359 |
| VA * | 18% | 12% | 1% | 69% | 762 |
| WI * | 19% | 10% | 29% | 42% | 362 |
| WY * | 41% | 34% | 6% | 20% | 424 |
| NCI-IDD Average | 35% | 31% | 4% | 31% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last Pap test (women 21 and older)




Table 7. Had a Pap test in the past three years (among women 21 and older)*Information may have been obtained through state records*

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home | |
|---------------------------------|----------------------------|------------|---------------------------|-------------------|-----------------------------|--------------------------------------|--------------------------------|------------|
| NV * | ↑ | 70% | 67 | n/a | 77% | n/a | n/a | n/a |
| GA | ↑ | 67% | 139 | n/a | 68% | n/a | 69% | n/a |
| PA | ↑ | 63% | 150 | n/a | 77% | n/a | 45% | n/a |
| CT * | ↑ | 63% | 195 | n/a | 76% | 71% | 43% | n/a |
| AL | ↑ | 62% | 120 | n/a | 69% | n/a | 33% | n/a |
| NJ | ↑ | 62% | 133 | n/a | 76% | n/a | 45% | n/a |
| MD | ↑ | 60% | 161 | n/a | 70% | 76% | 49% | n/a |
| NY | ↑ | 57% | 397 | n/a | 63% | 70% | 50% | n/a |
| UT | | 66% | 44 | n/a | n/a | n/a | n/a | n/a |
| KY | | 59% | 75 | n/a | n/a | n/a | 50% | n/a |
| IL | | 56% | 78 | n/a | 65% | n/a | 39% | n/a |
| NE | | 56% | 121 | n/a | 68% | n/a | 31% | 71% |
| SC | | 55% | 137 | n/a | 62% | 73% | 30% | n/a |
| VA | | 54% | 114 | n/a | 59% | n/a | 51% | 50% |
| MN | | 50% | 117 | n/a | 43% | n/a | 48% | n/a |
| IN | | 50% | 316 | 52% | 63% | 66% | 36% | n/a |
| MT | | 49% | 94 | n/a | 43% | 71% | 40% | n/a |
| NCI-IDD Average | | 49% | 6,997 | 57% | 59% | 66% | 36% | 42% |
| MO | | 48% | 116 | n/a | 55% | 53% | 32% | n/a |
| KS * | | 47% | 90 | n/a | 44% | 50% | n/a | n/a |
| AR | | 46% | 92 | n/a | n/a | 65% | 22% | n/a |
| FL | | 45% | 174 | n/a | 58% | 81% | 26% | n/a |
| OK | | 45% | 135 | n/a | 55% | 51% | 29% | n/a |
| OR * | | 45% | 122 | n/a | 42% | n/a | 39% | n/a |
| NC | | 43% | 130 | n/a | 64% | n/a | 28% | n/a |
| WI | | 43% | 79 | n/a | n/a | 59% | 39% | n/a |
| TX | | 42% | 144 | n/a | n/a | n/a | 37% | 33% |
| WY | | 42% | 130 | n/a | n/a | 62% | 19% | n/a |
| MI | | 40% | 121 | n/a | 55% | n/a | 26% | n/a |
| DE | | 40% | 117 | n/a | 40% | n/a | 41% | n/a |
| HI | | 38% | 88 | n/a | n/a | n/a | 32% | 45% |
| NH | ↓ | 28% | 109 | n/a | n/a | n/a | 12% | 38% |

Table note: States with more than 25% missing or don't know are marked with *; The following states had a low count of responses (N<20) and were not shown: DC

Table 7b. Last Pap test (among women 21 and older, includes “not applicable,” missing, and “don’t know” responses)

| State (names abbreviated) | More than 3 years ago | In past 3 years | Missing | Not applicable | Don't know | N |
|---------------------------|-----------------------|-----------------|-----------|----------------|------------|---------------|
| AL | 10% | 15% | 3% | 61% | 11% | 479 |
| AR | 12% | 10% | 1% | 59% | 18% | 417 |
| CT * | 12% | 20% | 1% | 60% | 7% | 602 |
| DE | 17% | 11% | 0% | 59% | 13% | 419 |
| DC * | 0% | 1% | 62% | 25% | 12% | 354 |
| FL | 18% | 15% | 2% | 61% | 4% | 532 |
| GA | 11% | 23% | 0% | 59% | 7% | 406 |
| HI | 16% | 9% | 7% | 59% | 9% | 351 |
| IL | 9% | 11% | 2% | 60% | 19% | 400 |
| IN | 15% | 15% | 1% | 62% | 9% | 1,080 |
| KS * | 12% | 11% | 18% | 51% | 9% | 393 |
| KY | 7% | 10% | 1% | 61% | 21% | 439 |
| MD | 12% | 18% | 0% | 58% | 11% | 533 |
| MI | 11% | 7% | 0% | 59% | 23% | 670 |
| MN | 14% | 15% | 1% | 62% | 7% | 403 |
| MO | 15% | 14% | 0% | 63% | 9% | 403 |
| MT | 13% | 13% | 4% | 58% | 12% | 365 |
| NE | 13% | 16% | 1% | 59% | 11% | 420 |
| NV * | 5% | 11% | 0% | 56% | 28% | 419 |
| NH | 20% | 8% | 2% | 59% | 12% | 399 |
| NJ | 11% | 18% | 2% | 58% | 10% | 460 |
| NY | 10% | 14% | 0% | 62% | 14% | 1,682 |
| NC | 12% | 9% | 1% | 65% | 13% | 594 |
| OK | 19% | 15% | 0% | 65% | 2% | 400 |
| OR * | 16% | 13% | 0% | 63% | 7% | 418 |
| PA | 8% | 13% | 4% | 59% | 15% | 718 |
| SC | 12% | 15% | 0% | 61% | 12% | 511 |
| TX | 20% | 15% | 3% | 61% | 2% | 420 |
| UT | 4% | 8% | 2% | 64% | 22% | 359 |
| VA | 7% | 8% | 0% | 63% | 21% | 762 |
| WI | 12% | 9% | 4% | 57% | 17% | 362 |
| WY | 18% | 13% | 2% | 59% | 8% | 424 |
| NCI-IDD Average | 14% | 13% | 2% | 61% | 11% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last mammogram (women 50 and older)




Table 8. Had a mammogram test in the past two years (among women age 50 and over)

Information may have been obtained through state records

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|--------------|---------------------------|-------------------|-----------------------------|--------------------------------------|--------------------------------|
| CT * ↑ | 87% | 83 | n/a | 88% | n/a | n/a | n/a |
| PA ↑ | 84% | 93 | n/a | 85% | n/a | n/a | n/a |
| MD | 82% | 55 | n/a | 88% | n/a | n/a | n/a |
| FL | 82% | 44 | n/a | n/a | n/a | n/a | n/a |
| NJ | 82% | 55 | n/a | 84% | n/a | n/a | n/a |
| OK | 79% | 57 | n/a | 80% | 76% | n/a | n/a |
| AL | 79% | 52 | n/a | 79% | n/a | n/a | n/a |
| IN | 78% | 137 | 84% | 71% | 85% | 73% | n/a |
| SC | 78% | 54 | n/a | 77% | n/a | n/a | n/a |
| OR * | 76% | 45 | n/a | n/a | n/a | n/a | n/a |
| GA | 75% | 69 | n/a | 75% | n/a | n/a | n/a |
| NE | 74% | 42 | n/a | 65% | n/a | n/a | n/a |
| AR | 72% | 36 | n/a | n/a | 73% | n/a | n/a |
| NCI-IDD Average | 72% | 2,605 | 80% | 72% | 78% | 62% | 73% |
| MI | 72% | 43 | n/a | 68% | n/a | n/a | n/a |
| IL | 71% | 35 | n/a | 67% | n/a | n/a | n/a |
| MN | 71% | 35 | n/a | 64% | n/a | n/a | n/a |
| NC | 70% | 44 | n/a | n/a | n/a | n/a | n/a |
| NY | 70% | 172 | n/a | 72% | n/a | n/a | n/a |
| MO | 66% | 38 | n/a | 62% | n/a | n/a | n/a |
| NH | 65% | 43 | n/a | n/a | n/a | n/a | 65% |
| MT | 65% | 51 | n/a | 63% | n/a | n/a | n/a |
| KY | 64% | 36 | n/a | n/a | n/a | n/a | n/a |
| KS | 61% | 31 | n/a | 55% | n/a | n/a | n/a |
| WY | 56% | 43 | n/a | n/a | 62% | n/a | n/a |
| VA | 55% | 44 | n/a | 58% | n/a | n/a | n/a |
| HI ↓ | 49% | 41 | n/a | n/a | n/a | n/a | 48% |
| WI ↓ | 43% | 28 | n/a | n/a | n/a | n/a | n/a |
| NV ↓ | 42% | 31 | n/a | 40% | n/a | n/a | n/a |
| DE ↓ | 35% | 60 | n/a | 33% | n/a | n/a | n/a |


Table note: States with more than 25% missing or don't know are marked with *; The following states had a low count of responses (N<20) and were not shown: DC, TX, UT

Table 8b. Last mammogram test (among women age 50 and over, includes “not applicable,” missing, and “don’t know”)

| State (names abbreviated) | More than 2 years | In past 2 years | Missing | Not applicable | Don't know | N |
|---------------------------|-------------------|-----------------|-----------|----------------|------------|---------------|
| AL | 2% | 9% | 1% | 82% | 6% | 479 |
| AR | 2% | 6% | 0% | 85% | 6% | 417 |
| CT * | 2% | 12% | 0% | 84% | 2% | 602 |
| DE | 9% | 5% | 0% | 84% | 1% | 419 |
| DC * | 0% | 1% | 62% | 28% | 9% | 354 |
| FL | 2% | 7% | 1% | 89% | 2% | 532 |
| GA | 4% | 13% | 0% | 80% | 3% | 406 |
| HI | 6% | 6% | 3% | 82% | 3% | 351 |
| IL | 3% | 6% | 1% | 88% | 3% | 400 |
| IN | 3% | 10% | 0% | 85% | 3% | 1,080 |
| KS | 3% | 5% | 18% | 72% | 3% | 393 |
| KY | 3% | 5% | 0% | 85% | 6% | 439 |
| MD | 2% | 8% | 0% | 87% | 3% | 533 |
| MI | 2% | 5% | 0% | 86% | 7% | 670 |
| MN | 2% | 6% | 0% | 89% | 2% | 403 |
| MO | 3% | 6% | 0% | 88% | 2% | 403 |
| MT | 5% | 9% | 1% | 82% | 3% | 365 |
| NE | 3% | 7% | 0% | 88% | 2% | 420 |
| NV | 4% | 3% | 0% | 85% | 7% | 419 |
| NH | 4% | 7% | 0% | 87% | 2% | 399 |
| NJ | 2% | 10% | 1% | 84% | 3% | 460 |
| NY | 3% | 7% | 0% | 88% | 2% | 1,682 |
| NC | 2% | 5% | 1% | 91% | 2% | 594 |
| OK | 3% | 11% | 0% | 86% | 0% | 400 |
| OR * | 3% | 8% | 0% | 88% | 1% | 418 |
| PA | 2% | 11% | 3% | 81% | 3% | 718 |
| SC | 2% | 8% | 0% | 86% | 3% | 511 |
| TX | 1% | 3% | 0% | 96% | 0% | 420 |
| UT | 1% | 1% | 1% | 92% | 5% | 359 |
| VA | 3% | 3% | 0% | 89% | 5% | 762 |
| WI | 4% | 3% | 1% | 86% | 6% | 362 |
| WY | 4% | 6% | 0% | 88% | 1% | 424 |
| NCI-IDD Average | 3% | 7% | 1% | 87% | 3% | 25,424 |

Table note: States with more than 25% missing or don't know are marked with *

Last colorectal cancer screening (among people 45 to 75)



Please note: Response options are not mutually exclusive.

Table 9. Last colorectal cancer screening (among people age 45 to 75)

Response options are not mutually exclusive. Information may have been obtained through state records; don't know included in denominator

| State (names abbreviated) | Colonoscopy within the past 10 years | Sigmoidoscopy within the past 5 years | Fecal Occult Blood Test (FOBT) or Fecal Immunochemical Test (FIT) in the past year | Never had exam or screening | Don't know | N |
|---------------------------------|---|---|--|--------------------------------------|---------------|--------------|
| AL | 23% | 5% | 9% | 22% | 43% | 220 |
| AR | 19% | 1% | 3% | 21% | 55% | 152 |
| CT * | 68% | 0% | 5% | 14% | 14% | 235 |
| DE * | 18% | 3% | 3% | 5% | 77% | 147 |
| DC * | 16% | 0% | 1% | 0% | 84% | 97 |
| FL | 36% | 0% | 5% | 38% | 20% | 161 |
| GA | 35% | 0% | 1% | 37% | 26% | 210 |
| HI * | 15% | 0% | 11% | 22% | 54% | 94 |
| IL | 35% | 1% | 10% | 17% | 37% | 154 |
| IN | 35% | 1% | 4% | 26% | 35% | 423 |
| KS * | 42% | 4% | 5% | 12% | 37% | 103 |
| KY | 27% | 1% | 5% | 12% | 55% | 151 |
| MD | 34% | 1% | 1% | 24% | 37% | 156 |
| MI | 19% | 3% | 3% | 11% | 68% | 219 |
| MN | 42% | 2% | 8% | 23% | 27% | 124 |
| MO | 37% | 3% | 11% | 18% | 30% | 128 |
| MT | 40% | 1% | 4% | 25% | 30% | 163 |
| NE | 48% | 0% | 16% | 8% | 28% | 158 |
| NV * | 11% | 0% | 0% | 7% | 82% | 133 |
| NH | 38% | 2% | 5% | 24% | 34% | 125 |
| NJ | 57% | 0% | 6% | 16% | 22% | 161 |
| NC | 31% | 1% | 10% | 15% | 42% | 153 |
| OK | 45% | 1% | 6% | 46% | 2% | 149 |
| OR * | 29% | 0% | 13% | 39% | 19% | 124 |
| PA * | 30% | 1% | 4% | 10% | 56% | 280 |
| SC | 33% | 0% | 2% | 18% | 48% | 202 |
| TX | 39% | 2% | 0% | 39% | 24% | 41 |
| UT | 22% | 0% | 4% | 11% | 61% | 98 |
| VA | 16% | 0% | 4% | 18% | 63% | 250 |
| WI * | 18% | 1% | 4% | 18% | 59% | 138 |
| WY | 41% | 10% | 0% | 31% | 24% | 131 |
| NCI-IDD Average | 32% | 1% | 9% | 23% | 36% | 7,758 |

*Table note: States with more than 25% missing or don't know are marked with *; NY is not included in this table due to data that could not be validated.*

Table 9b. Last colorectal cancer screening (among people age 45 to 75, includes “not applicable,” missing, and “don’t know” responses)

| State (names abbreviated) | Valid response | Missing | Not applicable | Don't know | N |
|---------------------------|----------------|------------|----------------|------------|---------------|
| AL | 26% | 1% | 52% | 20% | 477 |
| AR | 16% | 0% | 63% | 20% | 417 |
| CT * | 33% | 58% | 3% | 6% | 602 |
| DE * | 8% | 0% | 65% | 27% | 419 |
| DC * | 5% | 63% | 10% | 23% | 354 |
| FL | 25% | 2% | 67% | 6% | 532 |
| GA | 38% | 0% | 48% | 14% | 406 |
| HI * | 13% | 11% | 62% | 15% | 351 |
| IL | 24% | 3% | 59% | 15% | 400 |
| IN | 26% | 1% | 60% | 14% | 1,078 |
| KS * | 16% | 19% | 55% | 10% | 390 |
| KY | 15% | 1% | 65% | 19% | 439 |
| MD | 18% | 0% | 71% | 11% | 533 |
| MI | 10% | 2% | 65% | 22% | 669 |
| MN | 22% | 1% | 68% | 8% | 403 |
| MO | 22% | 0% | 68% | 10% | 403 |
| MT | 32% | 2% | 52% | 14% | 365 |
| NE | 27% | 0% | 62% | 11% | 420 |
| NV * | 6% | 0% | 68% | 26% | 416 |
| NH | 21% | 1% | 68% | 11% | 399 |
| NJ | 27% | 2% | 63% | 8% | 460 |
| NC | 15% | 1% | 73% | 11% | 589 |
| OK | 37% | 0% | 63% | 1% | 400 |
| OR * | 24% | 70% | 0% | 6% | 418 |
| PA * | 17% | 4% | 56% | 22% | 717 |
| SC | 21% | 0% | 60% | 19% | 511 |
| TX | 7% | 1% | 90% | 2% | 420 |
| UT | 11% | 1% | 71% | 17% | 357 |
| VA | 12% | 2% | 65% | 21% | 761 |
| WI * | 16% | 6% | 56% | 22% | 362 |
| WY | 23% | 3% | 67% | 8% | 424 |
| NCI-IDD Average | 20% | 22% | 46% | 11% | 23,722 |

Table note: States with more than 25% missing or don't know are marked with *; NY is not included in this table due to data that could not be validated.

Had flu vaccination in last 12 months




Table 10. Had a flu vaccine in the past 12 months

Information may have been obtained through state records

Key: Up arrow (↑) represents Significantly Above Average; down arrow (↓) represents Significantly Below Average

| State (names abbreviated) | Average within State | N | Institutional Settings | Group Settings | Own Home or Apartment | Parent's or Relative's Home | Foster Care or Host Home |
|---------------------------------|----------------------------|---------------|---------------------------|-------------------|-----------------------------|--------------------------------------|-----------------------------------|
| UT * ↑ | 89% | 170 | 100% | 96% | n/a | 74% | 80% |
| NE ↑ | 88% | 381 | 100% | 91% | 88% | 89% | 84% |
| CT ↑ | 87% | 557 | n/a | 94% | 91% | 72% | n/a |
| MT ↑ | 84% | 301 | n/a | 91% | 81% | 72% | n/a |
| IL * ↑ | 84% | 270 | n/a | 91% | n/a | 70% | n/a |
| DC * ↑ | 84% | 61 | n/a | n/a | 87% | n/a | n/a |
| AL ↑ | 83% | 417 | n/a | 89% | 82% | 62% | n/a |
| VA * ↑ | 82% | 414 | n/a | 92% | 83% | 71% | 79% |
| KS * ↑ | 81% | 253 | n/a | 93% | 69% | 61% | n/a |
| MN ↑ | 80% | 377 | n/a | 86% | 82% | 71% | n/a |
| GA ↑ | 79% | 380 | n/a | 87% | 76% | 73% | 78% |
| MO ↑ | 79% | 370 | n/a | 89% | 81% | 71% | n/a |
| SC ↑ | 77% | 412 | n/a | 89% | 56% | 67% | n/a |
| NJ ↑ | 77% | 405 | n/a | 87% | 81% | 66% | n/a |
| IN ↑ | 77% | 1,014 | 89% | 84% | 76% | 63% | n/a |
| NH | 77% | 318 | n/a | 94% | 71% | 67% | 79% |
| OK | 76% | 390 | n/a | 85% | 78% | 63% | n/a |
| WI * | 75% | 244 | n/a | 88% | 72% | 70% | n/a |
| NC * | 73% | 381 | 100% | 85% | 73% | 62% | 74% |
| MD | 72% | 461 | n/a | 90% | 74% | 63% | n/a |
| PA * | 72% | 404 | 95% | 83% | 63% | 50% | n/a |
| DE * | 72% | 204 | n/a | 66% | n/a | 86% | n/a |
| AR | 72% | 335 | n/a | 87% | 76% | 63% | n/a |
| KY * | 72% | 271 | n/a | 87% | 64% | 55% | 81% |
| MI * | 71% | 381 | n/a | 90% | 67% | 54% | n/a |
| NCI-IDD Average | 71% | 21,105 | 91% | 83% | 68% | 60% | 72% |
| TX | 66% | 383 | n/a | 79% | 61% | 63% | 66% |
| WY | 65% | 385 | n/a | n/a | 67% | 48% | n/a |
| NV * | 63% | 175 | n/a | 62% | 65% | 63% | n/a |
| OR ↓ | 61% | 391 | n/a | 77% | 47% | 59% | 62% |
| HI * ↓ | 60% | 178 | n/a | n/a | n/a | 43% | 76% |
| NY ↓ | 59% | 1,343 | n/a | 69% | 44% | 52% | n/a |
| FL ↓ | 57% | 495 | n/a | 69% | 61% | 47% | n/a |

Table note: States with more than 25% missing or don't know are marked with *

Table 10b. Had a flu vaccine in the past year (includes missing and “don’t know” responses)

| State (names abbreviated) | Flu vaccine in the past year | Not in the past year | Missing | Don't know | N |
|---------------------------|------------------------------|----------------------|-----------|------------|---------------|
| AL | 72% | 15% | 3% | 10% | 479 |
| AR | 58% | 23% | 1% | 19% | 417 |
| CT | 80% | 12% | 1% | 7% | 602 |
| DE * | 35% | 14% | 0% | 51% | 419 |
| DC * | 14% | 3% | 63% | 20% | 354 |
| FL | 53% | 40% | 2% | 5% | 532 |
| GA | 74% | 20% | 0% | 6% | 406 |
| HI * | 30% | 21% | 29% | 20% | 351 |
| IL * | 57% | 11% | 3% | 30% | 400 |
| IN | 72% | 22% | 0% | 6% | 1,080 |
| KS * | 52% | 12% | 19% | 16% | 393 |
| KY * | 44% | 18% | 0% | 38% | 439 |
| MD | 63% | 24% | 1% | 12% | 533 |
| MI * | 40% | 17% | 2% | 41% | 670 |
| MN | 74% | 19% | 4% | 3% | 403 |
| MO | 72% | 20% | 0% | 8% | 403 |
| MT | 70% | 13% | 3% | 14% | 365 |
| NE | 80% | 11% | 1% | 8% | 420 |
| NV * | 26% | 16% | 1% | 58% | 419 |
| NH | 61% | 19% | 2% | 19% | 399 |
| NJ | 68% | 20% | 5% | 7% | 460 |
| NY | 47% | 33% | 0% | 20% | 1,682 |
| NC * | 47% | 17% | 1% | 35% | 594 |
| OK | 74% | 24% | 0% | 3% | 400 |
| OR | 57% | 36% | 1% | 5% | 418 |
| PA * | 41% | 16% | 8% | 36% | 718 |
| SC | 62% | 18% | 0% | 19% | 511 |
| TX | 60% | 31% | 7% | 2% | 420 |
| UT * | 42% | 5% | 4% | 49% | 359 |
| VA * | 44% | 10% | 1% | 44% | 762 |
| WI * | 51% | 17% | 30% | 3% | 362 |
| WY | 59% | 32% | 6% | 4% | 424 |
| NCI-IDD Average | 58% | 24% | 4% | 14% | 25,424 |

In the past 12 months, have you gone to the emergency room for any reason (to get care for yourself)?





Table 11. Went to the emergency room for any reason to get care for themselves in the past 12 months

Proxy respondents were allowed for this question

| State (names abbreviated) | Yes | No | N |
|---------------------------|------------|------------|---------------|
| AL | 22% | 78% | 459 |
| AR | 20% | 80% | 412 |
| CT | 30% | 70% | 591 |
| DE | 28% | 72% | 406 |
| DC | 23% | 77% | 342 |
| FL | 18% | 82% | 522 |
| GA | 19% | 81% | 402 |
| HI | 22% | 78% | 346 |
| IL | 19% | 81% | 382 |
| IN | 32% | 68% | 1,055 |
| KS | 23% | 77% | 339 |
| KY | 17% | 83% | 422 |
| MD | 21% | 79% | 505 |
| MI | 23% | 77% | 660 |
| MN | 23% | 77% | 390 |
| MO | 23% | 77% | 393 |
| MT | 33% | 67% | 341 |
| NE | 25% | 75% | 395 |
| NV | 27% | 73% | 390 |
| NH | 28% | 72% | 392 |
| NJ | 25% | 75% | 446 |
| NY | 24% | 76% | 1,615 |
| NC | 23% | 77% | 584 |
| OK | 28% | 72% | 393 |
| OR | 25% | 75% | 408 |
| PA | 26% | 74% | 683 |
| SC | 25% | 75% | 503 |
| TX | 21% | 79% | 396 |
| UT | 30% | 70% | 317 |
| VA | 34% | 66% | 757 |
| WI | 32% | 68% | 346 |
| WY | 30% | 70% | 400 |
| NCI-IDD Average | 24% | 76% | 24,697 |

Reasons for unexpected hospitalization



Please note: Response options are not mutually exclusive.

Table 12. Had an unexpected hospital admission due to dehydration, bowel obstruction, seizure, aspiration, or GERD, in the past year

Information may have been obtained through state records. Response options are not mutually exclusive. Percentage shows “Of those with an unexpected hospital admission due to dehydration, bowel obstruction, seizure, aspiration, or GERD, in the past year, the percent that were hospitalized for each condition.” Information may have been obtained through state records.

| State (names abbreviated) | Dehydration | Bowel obstruction | Seizure | Aspiration | Gastroesophageal Reflux Disease (GERD) | N |
|---------------------------|-------------|-------------------|------------|------------|--|------------|
| AL | 4% | 13% | 58% | 4% | 33% | 24 |
| AR | 5% | 8% | 49% | 5% | 44% | 39 |
| CT | 17% | 14% | 48% | 17% | 14% | 29 |
| DE | 18% | 15% | 14% | 15% | 68% | 73 |
| FL | 23% | 9% | 59% | 14% | 5% | 22 |
| KS | 15% | 20% | 30% | 15% | 45% | 20 |
| KY | 15% | 23% | 19% | 15% | 35% | 26 |
| MI | 7% | 12% | 64% | 12% | 21% | 42 |
| MT | 11% | 11% | 37% | 26% | 33% | 27 |
| NH | 10% | 10% | 38% | 0% | 48% | 21 |
| NY | 10% | 10% | 59% | 11% | 23% | 92 |
| NC | 7% | 17% | 53% | 13% | 17% | 30 |
| OK | 35% | 35% | 22% | 22% | 13% | 23 |
| PA | 15% | 8% | 42% | 21% | 29% | 52 |
| SC | 20% | 11% | 39% | 4% | 39% | 46 |
| UT | 23% | 18% | 50% | 18% | 23% | 22 |
| VA | 8% | 23% | 46% | 31% | 12% | 26 |
| NCI-IDD Average | 13% | 14% | 47% | 13% | 26% | 778 |

Table note: States with more than 25% missing or don't know are marked with *, the following states had a low count of responses (N<20) and were not shown: DC, GA, HI, IL, IN, MD, MN, MO, NE, NV, NJ, OR, TX, WI, WY.

In the past 6 months, have you fallen and hurt yourself?




Table 13. Fell and hurt themselves in the past six months*Proxy respondents were allowed for this question*

| State (names abbreviated) | Fell and hurt themselves in past 6 months | Did not fall and hurt themselves in the past 6 months | N |
|---------------------------|---|---|---------------|
| AL | 11% | 89% | 463 |
| AR | 18% | 82% | 400 |
| CT | 13% | 87% | 593 |
| DE | 14% | 86% | 411 |
| DC | 12% | 88% | 343 |
| FL | 9% | 91% | 521 |
| GA | 10% | 90% | 403 |
| HI | 14% | 86% | 346 |
| IL | 9% | 91% | 385 |
| IN | 18% | 82% | 1,050 |
| KS | 18% | 82% | 346 |
| KY | 11% | 89% | 421 |
| MD | 11% | 89% | 510 |
| MI | 11% | 89% | 660 |
| MN | 13% | 87% | 390 |
| MO | 12% | 88% | 400 |
| MT | 24% | 76% | 343 |
| NE | 15% | 85% | 399 |
| NV | 19% | 81% | 406 |
| NH | 15% | 85% | 394 |
| NJ | 11% | 89% | 450 |
| NY | 12% | 88% | 1,617 |
| NC | 12% | 88% | 586 |
| OK | 19% | 81% | 392 |
| OR | 19% | 81% | 404 |
| PA | 13% | 87% | 674 |
| SC | 9% | 91% | 504 |
| TX | 12% | 88% | 394 |
| UT | 20% | 80% | 322 |
| VA | 11% | 89% | 756 |
| WI | 19% | 81% | 349 |
| WY | 21% | 79% | 402 |
| NCI-IDD Average | 13% | 87% | 24,747 |



Human Services
Research Institute

Human Services Research Institute (HSRI)

2336 Massachusetts Avenue

Cambridge, MA 02140



NASDDDS

National Association of State Directors
of

Developmental Disabilities Services

National Association of State Directors of Developmental Disabilities Services (NASDDDS)

301 N Fairfax Street, Suite 101

Alexandria, VA 22314

May 2024

Questions/Comments: nci@nationalcoreindicators.org